

Diet For Stage 3 Kidney Disease

Diet For Stage 3 Kidney Disease

Downloaded from evoluir.abar.org.br by Logan & Cassius

UNDERSTANDING THE ROLE OF DIET IN STAGE 3 KIDNEY DISEASE

Receiving a diagnosis of Stage 3 chronic kidney disease (CKD) can feel overwhelming, but it also marks a critical turning point where proactive management can significantly slow disease progression. The kidneys, which filter waste and excess fluids from the blood, are functioning at a moderately reduced level—typically between 30 and 59 percent of normal capacity. At this stage, the right dietary choices become a cornerstone of treatment.

A well-planned diet for Stage 3 kidney disease is not a one-size-fits-all prescription. Instead, it is a personalized approach designed to reduce the workload on the kidneys, maintain electrolyte balance, and prevent complications such as high blood pressure, bone disease, and fluid buildup. By making informed adjustments to what you eat, you can protect your kidney function, support overall health, and maintain a good quality of life.

This article will walk you through the essential principles of eating well with Stage 3 CKD. You will learn which nutrients require careful monitoring, which foods to emphasize, which to limit, and how to build a sustainable eating pattern that fits your lifestyle.

WHAT CHANGES IN STAGE 3 KIDNEY DISEASE?

In Stage 3, the kidneys have sustained significant damage, though they are still working. As a result, they become less efficient at removing certain waste products and maintaining the balance of minerals and fluids. Two key markers of kidney function are estimated glomerular filtration rate (eGFR), which falls between 30 and 59 in this stage, and the presence of protein in the urine. Your healthcare team, which typically includes a nephrologist and a renal dietitian, will use these numbers along with blood tests to guide your dietary plan.

It is important to recognize that symptoms may be mild or even absent at this stage. Some people experience fatigue, changes in urination, swelling in the hands or feet, or mild itching. Because the disease can progress silently, following a kidney-friendly diet early can help preserve remaining function longer.

KEY NUTRITIONAL GOALS FOR A DIET FOR STAGE 3 KIDNEY DISEASE

While every individual's needs vary, most dietary plans for Stage 3 CKD focus on five major areas: protein intake, sodium restriction, phosphorus management, potassium balance, and fluid control. Let us examine each of these in detail.

Protein: Quality Over Quantity

Protein is essential for muscle maintenance, immune function, and healing. However, when kidneys are damaged, the waste products of protein metabolism—such as urea and creatinine—can build up in the blood. For this reason, many people with Stage 3 kidney disease are advised to moderately reduce their protein intake. The goal is not to eliminate protein but to consume it in the right amount and from high-quality sources.

Typically, a renal dietitian might recommend between 0.6 and 0.8 grams of protein per kilogram of body weight per day, but this number can vary. Emphasis is placed on **high biological value proteins**—those that contain all essential amino acids—such as lean meats, poultry, fish, eggs, and dairy. Plant proteins like beans, lentils, and tofu can also be included but require careful portioning because they are often higher in potassium and phosphorus.

The key is to avoid excessive protein intake while still meeting your body's basic needs. This approach helps reduce the accumulation of nitrogenous wastes, making it easier for your kidneys to keep up.

Sodium: Keeping Blood Pressure in Check

High blood pressure is both a cause and a consequence of kidney disease. Limiting sodium is one of the most effective dietary strategies for managing blood pressure and reducing fluid retention. For most people with Stage 3 CKD, the recommendation is to consume less than 2,000 to 2,300 milligrams of sodium per day—roughly one teaspoon of salt.

This means cutting back on processed foods, canned soups, fast food, deli meats, salty snacks, and condiments like soy sauce and ketchup. Instead, flavor your meals with fresh or dried herbs, spices, garlic, lemon juice, and salt-free seasoning blends. Reading nutrition labels becomes a daily habit, as sodium hides in many unexpected places, including bread, cheese, and salad dressings.

By controlling sodium intake, you not only help your kidneys but also reduce the risk of cardiovascular disease, which is common in the CKD population.

Phosphorus: Protecting Your Bones and

FOODS TO EMPHASIZE IN A DIET FOR STAGE 3 KIDNEY DISEASE

Blood Vessels

In healthy kidneys, excess phosphorus is excreted easily. But as kidney function declines, phosphorus can accumulate in the blood. High phosphorus levels leach calcium from your bones, making them weak and brittle, and can also contribute to vascular calcification, which increases heart disease risk.

During Stage 3, phosphorus restrictions may not yet be drastic, but it is wise to start being mindful. Foods naturally high in phosphorus include dairy products, nuts, seeds, whole grains, cola drinks, and processed foods with phosphate additives. Many renal diets recommend limiting dairy to one serving per day and choosing unprocessed foods whenever possible.

Your doctor may also prescribe phosphate binders—medications taken with meals to prevent phosphorus absorption. If you take these, it is important to take them exactly as directed.

Potassium: A Delicate Balance

Potassium is a mineral that helps nerves and muscles function properly, including the heart. In Stage 3 kidney disease, potassium levels may remain normal for many people, though some begin to experience hyperkalemia (high potassium). Whether you need to restrict potassium depends entirely on your blood test results.

If your potassium is elevated, you will need to limit high-potassium foods such as bananas, oranges, potatoes, tomatoes, avocados, spinach, and beans. Lower-potassium alternatives include apples, berries, grapes, cauliflower, green beans, and white rice. Boiling certain vegetables can also help leach out some of their potassium content.

If potassium levels are normal, you typically do not need to restrict it—but awareness is still important because levels can change quickly if your kidney function declines further.

Fluids: Individualized Guidance

In Stage 3, fluid restrictions are not usually necessary unless you develop significant fluid retention or swelling. Some people may still be able to follow their natural thirst. However, if you have high blood pressure or edema, your healthcare provider may recommend moderating fluid intake. Thirst can be managed by sucking on ice chips, using sugar-free hard candies, or rinsing your mouth with water.

Fluids include not just water but also coffee, tea, juice, soup, and even foods that melt at room temperature like gelatin and ice cream. Always follow the specific advice given by your medical team.

Building a kidney-friendly plate means focusing on fresh, whole foods that are naturally low in sodium, phosphorus, and—when needed—potassium. Here are some excellent choices:

- **Fresh or frozen vegetables:** Cauliflower, cabbage, bell peppers, zucchini, green beans, lettuce, and cucumbers are generally lower in potassium.
- **Fresh fruits:** Apples, berries, grapes, cherries, pineapple, and watermelon are good options for most people.
- **Lean proteins:** Skinless chicken, turkey, fish, and eggs in controlled portions.
- **Grains:** White rice, pasta, bread, and crackers made from refined flour, as whole grains are higher in phosphorus.
- **Healthy fats:** Olive oil, avocado oil, and unsalted butter or margarine.
- **Beverages:** Water, clear sodas like lemon-lime (not cola), and unsweetened herbal teas (check with your dietitian first).

FOODS TO LIMIT OR AVOID

Equally important is knowing what to reduce or eliminate. The following foods often require caution:

- **Processed and cured meats:** Bacon, ham, sausage, hot dogs, and deli meats are extremely high in sodium and phosphorus additives.
- **Canned foods:** Canned soups, vegetables, and beans are usually loaded with sodium. If you use them, choose "no salt added" versions and rinse thoroughly.
- **Dairy products:** Milk, yogurt, and cheese are high in phosphorus. Stick to small portions or alternatives like unsweetened almond or rice milk (check for phosphate additives).
- **Dark-colored sodas:** Cola and root beer contain phosphorus additives that are easily absorbed.
- **High-potassium foods (if needed):** Bananas, oranges, tomatoes, potatoes, spinach, avocados, and dried fruits.
- **Nuts, seeds, and legumes:** These are nutrient-dense but also high in both potassium and phosphorus. Use in very small amounts if permitted.
- **Salt and salty condiments:** Soy sauce, Worcestershire sauce, ketchup, pickles, and olives.

SAMPLE MEAL IDEAS FOR STAGE 3 KIDNEY DISEASE

Putting theory into practice can be the hardest part. Here are a few simple meal ideas that align with the dietary goals for Stage 3 CKD. Remember to adjust based on your individual lab values.

Breakfast

Scrambled eggs (using one egg) with a side of sautéed bell peppers and onions (cooked in olive oil). Serve with a small apple and a slice of white toast with unsalted butter. Drink water or a cup of light

herbal tea.

Lunch

Grilled chicken breast (about 3 ounces) over a bed of mixed greens with cucumber slices, shredded carrots, and a vinaigrette made from olive oil, vinegar, and dried herbs. Add a side of steamed white rice and a handful of grapes.

Dinner

Baked salmon (3 to 4 ounces) seasoned with lemon, dill, and garlic powder. Serve with roasted green beans and a small baked potato that has been boiled (to leach potassium) or limited to half a potato if potassium is a concern. A side of cauliflower rice makes a nice addition.

Snack

A small bowl of fresh blueberries or a few unsalted rice cakes with a thin layer of unsalted butter or sugar-free jam. Avoid high-potassium snacks like bananas or mixed nuts.

LIFESTYLE TIPS FOR MANAGING KIDNEY HEALTH

Diet is only one part of the picture. To get the most out of your renal diet, consider these additional strategies:

- **Work with a renal dietitian:** A professional can help you tailor the diet to your specific lab results, preferences, and lifestyle. This is the single most effective step you can take.

- **Monitor your blood pressure regularly:** Keeping blood pressure below 130/80 is a common target. Diet, exercise, and medication all play a role.
- **Stay active:** Gentle exercise such as walking, swimming, or yoga helps control weight, blood pressure, and overall well-being. Always check with your doctor before starting a new routine.
- **Avoid NSAIDs:** Over-the-counter pain relievers like ibuprofen and naproxen can damage the kidneys further. Use acetaminophen instead, but only as directed.
- **Stay on top of lab work:** Your eGFR, creatinine, potassium, phosphorus, and other markers will guide dietary adjustments. Never make major changes without consulting your healthcare team.
- **Manage other conditions:** Diabetes and heart disease are common in people with kidney disease. Controlling blood sugar and cholesterol is essential.

COMMON QUESTIONS ABOUT DIET FOR STAGE 3 KIDNEY DISEASE

Can I eat eggs on a kidney diet?

Yes, eggs are an excellent source of high-quality protein and are lower in potassium and phosphorus than many other protein sources. One egg per day is generally acceptable, but always follow your dietitian's portion guidance.

Is it safe to drink coffee or tea?

In moderation, yes. Plain coffee and tea are low in sodium and phosphorus. However, they do contain some potassium. If your potassium is normal, one or two cups per day are usually fine. Avoid adding milk or creamer,