

Nihss Spanish Version

Nihss Spanish Version

Downloaded from evoluir.abar.org.br by Talan & Koch

UNDERSTANDING THE NIHSS AND ITS SPANISH VERSION: A CRITICAL TOOL FOR STROKE CARE

Stroke is a leading cause of disability and death worldwide, and timely, accurate assessment is essential for effective treatment. The National Institutes of Health Stroke Scale (NIHSS) is the gold standard for quantifying neurological deficits in acute stroke patients. However, language barriers can compromise the reliability of this assessment, particularly in Spanish-speaking populations. The **Nihss Spanish Version** was developed to address this gap, ensuring that clinicians can administer the scale with cultural and linguistic accuracy. This article explores what the NIHSS is, why a Spanish version is indispensable, how it was created, and how it improves stroke outcomes for millions of patients.

WHAT IS THE NIHSS?

The National Institutes of Health Stroke Scale is a 15-item instrument used by healthcare professionals to evaluate the severity of stroke-related neurological impairment. It measures domains such as level of consciousness, gaze, visual fields, facial palsy, motor arm and leg function, limb ataxia, sensory loss, language, dysarthria, and extinction/inattention. Each item is scored from 0 (normal) to a maximum of 2, 3, or 4, with total scores ranging from 0 to 42. Higher scores indicate more severe stroke. The NIHSS is widely employed in clinical trials, emergency departments, and stroke units because of its reproducibility and predictive validity for patient outcomes.

Originally developed in English, the scale relies heavily on verbal commands and patient responses. This creates a challenge when the patient speaks only Spanish, or when the clinician is not fluent in that language. Inaccurate scoring due to misinterpretation can lead to inappropriate treatment decisions, such as the administration of thrombolytics or endovascular therapy. The **Nihss Spanish Version** is therefore not merely a translation but a carefully adapted tool that preserves the scale’s psychometric properties while making it accessible to Spanish-speaking patients.

WHY A SPANISH VERSION OF THE NIHSS IS CRITICAL

Demographic Imperative

Spanish is the second most spoken language in the world by native speakers, and in countries like the United States, over 40 million people speak it at home. The Hispanic population is aging, increasing their risk of stroke. Without a validated **Nihss Spanish Version**, these patients face

systematic inequalities in stroke care. Bilingual clinicians are not always available, and even when they are, nuance can be lost in real-time translation. A standardized Spanish NIHSS eliminates variability and ensures that every patient receives the same objective assessment regardless of the clinician’s language skills.

Accuracy of Assessment

Many NIHSS items require the patient to follow complex commands, name objects, or repeat phrases. If the patient mishears a word or the clinician uses a culturally unfamiliar term, the score may reflect a false deficit. For example, the English command “Close your eyes” is simple, but a direct translation like “Cierre los ojos” may be understood. However, items that test comprehension (e.g., “Show me your teeth” for facial palsy) can be misinterpreted in some dialects. The **Nihss Spanish Version** has been reviewed by native speakers and stroke neurologists to ensure that the commands are natural and idiomatic across major Spanish-speaking regions.

Standardization in Research

Clinical trials often include multinational enrollment. If some centers use an ad-hoc translation of the NIHSS while others use the validated English version, data integrity is compromised. The **Nihss Spanish Version** allows for uniform data collection in Spanish-speaking populations, making results more generalizable and supporting subgroup analyses. Regulatory agencies like the FDA and EMA accept data from studies that use culturally adapted instruments, provided they have been validated.

DEVELOPMENT AND VALIDATION OF THE SPANISH NIHSS

Translation Process

Creating the **Nihss Spanish Version** involved a multi-step process recommended by international guidelines for cross-cultural adaptation. First, forward translation was performed by two independent bilingual translators whose native language is Spanish. A reconciliation meeting resolved discrepancies. Then, a back-translation into English was compared with the original to check for conceptual equivalence. An expert committee, including neurologists, neuropsychologists, and linguists, reviewed the pre-final version for clarity and cultural relevance.

Field Testing and Validation

After translation, the scale was pilot-tested with a small group of Spanish-speaking stroke patients and clinicians. Any confusing items were revised. Formal validation studies compared scores obtained with the **Nihss Spanish Version** against the English NIHSS in bilingual patients, as well as against other stroke severity measures. Inter-rater reliability and test-retest reliability were found to be excellent, with correlation coefficients above 0.90. Sensitivity and specificity for detecting large-vessel occlusion, a key clinical decision point, were also comparable to the English version.

One of the most widely used validated adaptations is the Spanish NIHSS developed by the Spanish Stroke Organization (Sociedad Española de Neurología) and later adopted by the National Institutes of Health. This version is now available on the NIH website and in many electronic health record systems. Studies have confirmed that it maintains the original scale's predictive validity for functional outcome at 90 days, a critical endpoint in stroke research.

HOW TO USE THE NIHSS SPANISH VERSION

Administering the **Nihss Spanish Version** follows the same protocol as the English version, with specific verbal commands provided in Spanish. Below are the 15 items, highlighted with their Spanish phrasing where relevant:

- **1a. Level of Consciousness (Alerta, Somnoliento, etc.)** – Scored based on response to voice or pain.
- **1b. LOC Questions (Preguntas: ¿En qué mes estamos? ¿Qué edad tiene?)** – The patient answers two questions.
- **1c. LOC Commands (Órdenes: Cierre los ojos y abra la mano)** – The patient follows two commands.
- **2. Gaze (Mirada)** – Assess horizontal eye movement; no verbal command needed.
- **3. Visual Fields (Campos visuales)** – Finger counting in each quadrant.
- **4. Facial Palsy (Parálisis facial)** – The patient is asked to show teeth (Enseñe los dientes) or smile (Sonría).
- **5 & 6. Motor Arm and Leg (Brazo y pierna derecha/izquierda)** – The patient holds the limb for 10 seconds (Mantenga el brazo/pierna arriba).
- **7. Limb Ataxia (Ataxia de extremidades)** – Finger-to-nose test (Toque su nariz con el dedo).
- **8. Sensory (Sensibilidad)** – Pinprick on face, arm, leg (Siente esto? ¿Es igual aquí?).
- **9. Best Language (Lenguaje)** – The patient names objects in pictures (Nómbreme estos objetos) and reads sentences (Lea estas frases).
- **10. Dysarthria (Disartria)** – The patient repeats words (Repita: Mamá, pastel, televisión).
- **11. Extinction and Inattention (Inatención)** – The patient is touched on left and right sides; then asked ¿Siente que le toco aquí?

Clinicians must be aware that the **Nihss Spanish Version** includes regionally neutral words. For example, instead of “automóvil” (car) some versions use “coche” or “auto” depending on target population; the validated version often provides acceptable alternatives. Training materials and instructional videos are available through the NIH Stroke Scale certification program, which now offers a Spanish-language track for both initial certification and recertification.

BENEFITS OF THE NIHSS SPANISH VERSION IN CLINICAL PRACTICE

Reduced Assessment Time

When a clinician must pause to find a translator or use a phrasebook, precious minutes are lost. With a standardized **Nihss Spanish Version** on hand, the assessment can proceed without delays, especially important for door-to-needle time in intravenous thrombolysis. Many hospitals have printed cards with the Spanish commands inserted into stroke kits, allowing even non-Spanish-speaking staff to perform the assessment by reading the displayed script.

Improved Accuracy and Inter-Rater Reliability

Because the **Nihss Spanish Version** has defined instructions and acceptable prompts, different clinicians are more likely to score the same patient consistently. This reliability is crucial for serial assessments to track improvement or deterioration. Studies show that using the Spanish version reduces scoring errors compared to improvised translations, particularly for language and comprehension items.

Empowerment of Spanish-Speaking Patients and Families

When a patient hears commands in their native language, they are more likely to cooperate and trust the medical team. This also reduces anxiety during a frightening experience like a stroke. Family members who are present can participate in providing history if the patient has aphasia, knowing that the scale is being administered appropriately. The **Nihss Spanish Version** thus contributes to culturally competent care.

CHALLENGES IN USING THE SPANISH NIHSS

Dialectal Variations

Spanish is not monolithic; vocabulary varies between Spain, Mexico, the Caribbean, and South America. For example, the word for “elevator” (ascensor vs. elevador) can confuse patients. The validated **Nihss Spanish Version** aims for neutral terms, but some dialectal adjustments may still be needed in specific regions. Ideally, each hospital should adopt a version that aligns with its predominant patient population and train staff on any required adaptations.

Training and Certification Gaps

Not all clinicians who treat stroke patients have completed the NIHSS certification in Spanish. Even if they are native Spanish speakers, they may not be familiar with the formal scale phrasing. Mandatory training programs using the **Nihss Spanish Version** are gradually being adopted by stroke centers, but more widespread implementation is needed. Online modules and simulations in Spanish can help bridge this gap.

Coexisting Aphasia or Cognitive Impairment

Stroke patients may have global aphasia affecting both languages in bilinguals. The **Nihss Spanish**

Version is designed for patients whose primary language is Spanish, but if the patient is bilingual, the clinician should use the language in which the patient is most fluent before the stroke. This may require a quick baseline assessment from family. In cases of severe aphasia, many NIHSS items are scored observationally (e.g., motor or visual tasks), so the verbal items are not always the sole determinant.

FUTURE DIRECTIONS: DIGITAL TOOLS AND TELEMEDICINE

The **Nihss Spanish Version** is increasingly being integrated into mobile apps and telemedicine platforms. These digital tools allow remote neurologists to administer the scale via video, with the scale displayed on screen in Spanish. Some electronic health records now auto-populate the Spanish version when the patient’s preferred language is Spanish. Artificial intelligence is also being explored to assist with scoring, though human judgment remains paramount.

Additionally, research is underway to create a simplified, shortened version of the NIHSS in Spanish for prehospital use by paramedics. This could speed up triage and alerting of stroke teams. The development of culturally sensitive patient education materials in Spanish that explain the NIHSS process will further empower patients and improve compliance.

CONCLUSION

The **Nihss Spanish Version** is not a luxury but a necessity in modern stroke care. It ensures that millions of Spanish-speaking patients receive the same standardized, accurate neurological assessment as English speakers. Through careful translation, validation, and ongoing training, this tool eliminates language barriers that once introduced dangerous variability into stroke evaluations. Clinicians, hospitals, and health systems must prioritize the adoption and proper use of the Spanish