
How To Grow Magic Mushrooms

How To Grow Magic Mushrooms

Downloaded from evoluir.abar.org.br by
Oscar & Griffith

INTRODUCTION TO CULTIVATING PSILOCYBIN MUSHROOMS

The interest in psychedelic compounds has surged in recent years, with a growing body of research highlighting the therapeutic potential of psilocybin for conditions such as treatment-resistant depression, anxiety, and PTSD. As a result, many individuals are curious about the process behind these fungi. However, before exploring the details of **how to grow magic mushrooms**, it is imperative to understand the legal landscape. In most countries, including the United States (at the federal level), the cultivation, possession, and consumption of psilocybin-containing mushrooms remain illegal. This article is intended for educational and informational purposes only, focusing on the mycological science behind the growth process. We strongly advise readers to check their local laws and regulations before considering any form of cultivation.

Growing mushrooms, in general, is a fascinating exercise in patience and microbiology. Unlike plants, fungi do not rely on photosynthesis. Instead, they require a sterile environment, a nutrient-rich substrate, and precise humidity and temperature

controls. The journey from spore to mature mushroom is a delicate dance of biology, requiring the cultivator to act as both a scientist and a caretaker. This guide will walk you through the foundational principles of indoor cultivation, emphasizing the importance of cleanliness and attention to detail.

UNDERSTANDING THE MAGIC MUSHROOM LIFECYCLE

To successfully cultivate any mushroom, you must first understand its lifecycle. Magic mushrooms, belonging primarily to the genus *Psilocybe*, follow a specific biological rhythm. The process begins with **spores**, which are microscopic reproductive units. Think of spores as seeds, though they are far more vulnerable to contamination. When spores land on a suitable medium, they germinate into a network of thread-like cells called **mycelium**. The mycelium colonizes the substrate, breaking down nutrients and preparing the environment for fruiting. Once the mycelium has fully colonized the substrate and environmental conditions are triggered (such as a drop in temperature or an increase in fresh air), **primordia** (tiny pinheads) form. These pins grow into mature **fruiting bodies**, the mushrooms themselves, which eventually release spores to restart the cycle.

Understanding this lifecycle is crucial because every step in the cultivation process is designed to facilitate one of these stages.

When you learn **how to grow magic mushrooms**, you are essentially learning how to create a controlled micro-environment that mimics the fungus's natural habitat while excluding competing organisms like bacteria and mold.

The Importance of Sterile Technique

Before purchasing any supplies, you must internalize one golden rule of mycology: **sterility**. Contamination is the single greatest enemy of the mushroom cultivator. A single bacterial spore or mold spore landing on your nutrient-rich substrate can outcompete the slow-growing psilocybe mycelium, ruining an entire batch. Successful cultivation relies on a rigorous sterile workflow, often referred to as the "clean technique." This involves using a pressure cooker to sterilize substrates, working inside a still-air box (SAB) or in front of a laminar flow hood, and using 70% isopropyl alcohol to disinfect all surfaces and tools. Without a firm commitment to cleanliness, your efforts are likely to fail.

ESSENTIAL SUPPLIES FOR CULTIVATION

Learning **how to grow magic mushrooms** begins with gathering the right equipment. While there are advanced methods, beginners often start with the "PF Tek" (Psilocybe Fanaticus Technique), which uses brown rice flour and vermiculite. Here is a list of fundamental supplies:

- **Spore Syringe:** This is a sterile syringe containing

mushroom spores suspended in water. It is used to inoculate your substrate. Note that spore syringes are often sold for microscopy purposes only.

- **Substrate:** The food source for the mycelium. Common choices include brown rice flour, vermiculite, and water, or more advanced options like rye grain or wild bird seed.
- **Glass Jars:** Wide-mouth half-pint or quart jars are used to hold the substrate during colonization. They must be able to withstand high heat in a pressure cooker.
- **Pressure Cooker:** Essential for sterilizing the substrate. A stovetop pressure cooker capable of reaching 15 PSI is the industry standard.
- **Still-Air Box (SAB):** A large plastic tote with holes cut for your arms. This creates a still environment where airborne contaminants are less likely to fall onto your work.
- **Mist Sprayer and Vermiculite:** Used for maintaining humidity in the fruiting chamber (a transparent plastic tub).
- **Lighter and Isopropyl Alcohol:** For flame-sterilizing the needle of the syringe and disinfecting surfaces.

STEP-BY-STEP GUIDE: FROM SPORE TO HARVEST

Now, let us break down the process into actionable steps. This guide assumes you are using a standard PF Tek method with a shotgun fruiting chamber (SGFC).

Step 1: Preparing the Substrate (Jars)

The substrate provides the nutrients and physical structure for the mycelium. For a PF Tek, you will mix **brown rice flour** with **vermiculite** and water. Vermiculite holds moisture and provides air pockets, while the rice flour is the nutrient source. Mix 2 parts vermiculite, 1 part brown rice flour, and 1 part water by volume. Pack this mixture loosely into half-pint wide-mouth jars, leaving about half an inch of headspace. Wipe the rims clean, and cover the jar openings with a layer of dry vermiculite (the "dry vermiculite layer"). This acts as a filter. Secure the lids, but leave them slightly loose to allow for gas exchange. Finally, cover the jar tops with a sheet of aluminum foil to prevent water from entering during sterilization.

Step 2: Sterilization

Place the jars in your pressure cooker. Add water according to your cooker's instructions (usually about 2-3 inches deep). Heat the cooker until it reaches 15 PSI and maintain that pressure for 60 minutes. This process kills any latent contaminants inside the substrate. Allow the cooker to cool down naturally. Do not rush this; forced cooling can suck contaminated air into the jars. Once the jars are cool to the touch (often left to sit overnight), they are ready for inoculation.

Step 3: Inoculation

This is where you introduce the spores to the sterile substrate. Work inside your still-air box. First, thoroughly wipe down the inside of the SAB and your hands with 70% isopropyl alcohol. Place your spore syringe and cooled jars inside. Flame-sterilize the syringe needle until it glows red, then let it cool for a few seconds. Hold the jar at an angle and inject about 0.5 to 1 cc of spore solution through the dry vermiculite layer. Inject into the substrate, avoiding the glass. Repeat for all jars. Immediately cover the injection hole with micropore tape. Clean up and store the jars in a dark, warm place (75-80°F or 24-27°C) for colonization.

Step 4: Colonization (Mycelium Growth)

Over the next 1 to 3 weeks, you will watch the white, fuzzy mycelium spread through the brown rice flour substrate. This is the most satisfying visual stage. The goal is to avoid "contams" (contamination). If you see green, black, or red patches (trichoderma or other molds), or if the jar smells sour (bacterial fermentation), remove that jar immediately and dispose of it away from your grow area. A healthy jar will smell earthy, like fresh mushrooms. Once the mycelium has fully colonized the substrate and the jar looks solid white, you can proceed to the fruiting stage. It is beneficial to wait an additional 5 to 7 days

after full colonization to allow the mycelium to "consolidate" and build strength.

Step 5: Fruiting (The Shotgun Fruiting Chamber)

Fruiting requires a change in environment. You need a chamber that provides high humidity and fresh air exchange. A "Shotgun Fruiting Chamber" (SGFC) is the standard. This is a clear plastic tote with 1/4-inch holes drilled every 2 inches on all six sides. Fill the bottom of the SGFC with 2-3 inches of moistened perlite. Perlite holds water and releases humidity. Once the jars are fully colonized, remove the mushroom "cake" from each jar. You can do this by gently tapping or using a clean, sterilized knife to loosen the cake. Rinse the cakes under cool, distilled water to remove the dry vermiculite layer. Soak the cakes in a bowl of distilled water for 12-24 hours (this is called "dunking" and rehydrates the mycelium for fruiting).

After dunking, roll the cakes in dry vermiculite; this helps retain moisture. Place the cakes on the metal mesh (or wire rack) inside the SGFC, resting them on the moist perlite. The holes in the chamber provide passive fresh air exchange. Mist the walls of the chamber and the perlite with water several times a day to keep the humidity near 99%. Fan the chamber with the lid to induce evaporation, which triggers pin formation.

Step 6: Harvesting

Within 5 to 14 days, you will see tiny white bumps (primordia) forming on the cakes. These grow into "pins" and eventually mature mushrooms. Harvest your mushrooms just as the **veil** (the thin membrane covering the gills underneath the cap) begins to tear but before it fully opens and drops spores. This is the peak of potency and visual appeal. Grip the mushroom at the base and gently twist and pull to remove it. If you cut it, the stump left behind can rot and cause contamination. Harvest all mushrooms promptly. After the first flush, you can re-dunk the cakes for another 12 hours and place them back in the SGFC for a second flush. Most growers get 2-3 flushes per cake.

DRYING AND STORAGE

Fresh mushrooms contain a lot of water and will rot quickly. To preserve potency for future use (therapeutic, microdosing, or research), you must dry them thoroughly. A **food dehydrator** set to a low temperature (under 150°F or 65°C) is the most effective method. Run the dehydrator until the mushrooms are "cracker dry," meaning they snap cleanly when bent. You can also use a desiccant like DampRid in a sealed container for a slower, non-heat method. Once dried, store the mushrooms in an airtight glass jar with a desiccant packet and keep them in a cool, dark place. Properly stored, they can retain potency for years.

Understanding **how to grow magic mushrooms** is not just about the final product; it is a deep dive into microbiology, mycology, and patience. The skills you learn—sterile technique,

environmental control, and observation—are applicable to growing any type of edible or medicinal mushroom, such as shiitake or lion's mane.

COMMON PITFALLS AND TROUBLESHOOTING

Even experienced cultivators face challenges. Here are some common issues you might encounter:

- **Contamination:** The most common problem. Usually caused by poor sterilization or sloppy technique. Prevention is your only viable weapon; once mold sets in, the batch is lost.
- **Slow Growth:** Often due to temperatures that are too low. Mycelium is slow below 70°F. Conversely, temperatures above 85°F can kill the mycelium and invite bacteria.
- **No Pins (Fruiting bodies):** This usually relates to poor fresh air exchange or lack of evaporation. Increase fanning and ensure your SGFC has adequate holes.
- **Thin, Leggy Mushrooms:** These are "anaerobic" or "etiolated" mushrooms stretching for air. They indicate that your CO2 levels are too high. Increase fresh air exchange dramatically.
- **Aborts (Small mushrooms that stop growing):** These can result