
Sterile Products And Aseptic Techniques

*Sterile Products And
Aseptic Techniques*

*Downloaded from
evoluir.abar.org.br by
Hooper & Autumn*

STERILE PRODUCTS AND ASEPTIC TECHNIQUES: THE FOUNDATION OF SAFE PHARMACEUTICAL COMPOUNDING

In the world of modern healthcare, the safety and efficacy of medications are paramount. At the heart of this commitment to patient well-being lies a critical discipline: the preparation of

sterile products and the practice of aseptic techniques. For patients who are immunocompromised, critically ill, or undergoing complex treatments like chemotherapy or parenteral nutrition, the difference between a properly prepared sterile product and one that has been contaminated can be a matter of life or death. Understanding the intricate relationship between sterile products and aseptic techniques is essential not only for pharmacy

professionals but for anyone involved in the healthcare delivery chain.

This article offers a deep dive into the world of sterile compounding, exploring the principles, procedures, and environmental controls that ensure products remain free from microbial contamination. By demystifying these complex processes, we aim to underscore why aseptic technique is not just a checklist of steps, but a mindset of meticulous attention to detail and unwavering commitment to patient safety.

WHAT ARE STERILE PRODUCTS?

Sterile products are pharmaceutical preparations that are free from all viable microorganisms, including bacteria, fungi, viruses, and their spores. These

products are typically administered via routes that bypass the body's natural defensive barriers, such as the skin or mucous membranes. Common routes of administration for sterile products include intravenous, intramuscular, intrathecal, epidural, and ophthalmic delivery.

The category of sterile products encompasses a wide range of medications, including:

- **Intravenous (IV) solutions** such as saline, dextrose, and lactated Ringer's solution.
- **Injectable medications** like antibiotics, analgesics, and insulin.
- **Chemotherapy drugs** that require precise dosing and absolute sterility.

- **Parenteral nutrition** (TPN) solutions that provide complete nutritional support intravenously.
- **Ophthalmic preparations** such as eye drops and ointments for sterile ophthalmic use.
- **Irrigation solutions** used in surgical procedures.

The critical nature of these products demands that they be prepared and handled under strict conditions that prevent the introduction of any microbial contaminants. A single microorganism introduced into the bloodstream can lead to a life-threatening infection, sepsis, or even death, particularly in a vulnerable patient population.

The Difference Between Sterile and Non-Sterile Compounding

It is important to distinguish between sterile and non-sterile compounding. Non-sterile compounding involves the preparation of medications that are administered orally, topically, or via other routes where the body's natural defenses can manage low levels of microbial presence. Sterile compounding, on the other hand, demands a much higher level of

environmental control, specialized training, and rigorous aseptic technique because the product will be introduced directly into sterile body sites such as the bloodstream or the central nervous system.

THE FUNDAMENTALS OF ASEPTIC TECHNIQUES

Aseptic technique is a set of specific practices and procedures performed with the goal of preventing contamination from pathogens. It involves applying the strictest rules to minimize the risk of infection and is a cornerstone of sterile product preparation. The term "aseptic" literally means "without sepsis," or without the presence of disease-causing microorganisms.

Aseptic technique is not limited to the

moment of compounding. It encompasses the entire workflow, from the moment a practitioner enters the compounding area to the final check before a product is released for administration. It is a comprehensive system that includes personal hygiene, environmental controls, equipment maintenance, and procedural discipline.

Why Aseptic Technique Matters for

Sterile Products

The connection between sterile products and aseptic techniques is inseparable. Without proper aseptic technique, even the cleanest environment and the most sterile starting materials can become contaminated during the compounding process. The human hand, for example, carries thousands of microorganisms. A single improper touch on a critical site, such as the tip of a syringe or the surface of a vial stopper, can introduce contaminants that proliferate in the nutrient-rich environment of an IV solution.

Consider the case of a patient receiving total parenteral nutrition. The solution contains glucose, amino acids, lipids,

and electrolytes—an excellent growth medium for bacteria and fungi. If a contaminant is introduced during compounding, it can multiply rapidly within the solution before infusion, leading to a devastating bloodstream infection. This reality underscores why every action in the sterile compounding suite must be deliberate, careful, and guided by the principles of aseptic technique.

CORE PRINCIPLES OF ASEPTIC TECHNIQUE FOR STERILE PRODUCT COMPOUNDING

The practice of aseptic technique is built upon several foundational principles that must be consistently applied. These principles are designed to create and maintain a sterile field, protect both the

product and the patient, and ensure the integrity of the final sterile product.

1. Hand Hygiene and Personal Garbing

Proper hand hygiene is the single most important step in preventing contamination. Before entering the sterile compounding area, personnel must perform a thorough hand wash using an antimicrobial soap, paying special attention to the fingernails and between the fingers. The hands must be dried with sterile lint-free wipes to prevent moisture from compromising

sterile barriers.

Following hand hygiene, complete garbing is required. This typically includes:

- A sterile, low-lint gown that covers personal clothing and exposed skin.
- A head cover that contains all hair.
- A face mask to prevent droplet contamination from speaking, coughing, or sneezing.
- Shoe covers to prevent tracking contaminants from outside the cleanroom.
- Sterile, powder-free gloves, which are often wiped with 70% isopropyl alcohol immediately before handling sterile products.

2. Environmental Controls and Cleanroom Standards

Sterile compounding must take place within a controlled environment that meets stringent standards for air quality, temperature, and pressure. These environments are typically classified according to International Organization for Standardization (ISO) classifications. For example, ISO Class 5 environments have fewer than 3,520 particles (0.5 microns or larger) per cubic meter of air.

Key components of the sterile compounding environment include:

- **Laminar airflow workstations (LAFW):** These provide an ISO Class 5 environment by directing HEPA-filtered air across the work surface in a unidirectional flow, sweeping away airborne contaminants.
- **Biological safety cabinets (BSC):** Used for hazardous drugs like chemotherapy agents, these cabinets provide both product protection and personnel protection through HEPA filtration and directional airflow.
- **Cleanroom anteroom:** A buffer zone between the outside environment and the sterile

compounding area, where garbing and hand hygiene are performed.

- **Positive air pressure:**

Cleanrooms are maintained at positive pressure relative to adjacent areas to prevent unfiltered air from entering the sterile field.

3. Workflow and Material Flow

Proper workflow is essential to maintaining the integrity of sterile products. The concept of "first air" refers to the clean, HEPA-filtered air that first contacts the sterile product and critical sites. All manipulations must be

performed within this first air zone, and nothing should come between the HEPA filter and the work area that could disrupt the airflow.

Materials should be organized efficiently before compounding begins. All necessary components—syringes, needles, vials, IV bags, and diluents—should be gathered and inspected for defects, expiration dates, and particulate matter. The work surface must be disinfected with 70% isopropyl alcohol, and all items introduced into the workspace should be wiped down with alcohol to remove surface contaminants.

4. Critical Site Management

A critical site is any area on a sterile product or device that comes into direct contact with the sterile product or the patient. Examples include the tip of a syringe, the needle hub, the diaphragm of a vial stopper, and the injection port of an IV bag. These sites must never be touched by non-sterile objects, including ungloved hands, unprotected surfaces, or airborne contaminants.

Management of critical sites involves:

- Disinfecting vial stoppers and injection ports with 70% isopropyl alcohol before each use.

- Allowing the alcohol to dry completely, which is critical for achieving effective microbial kill.
- Using aseptic connections that maintain sterility at every junction.
- Never leaving a critical site exposed to the open environment longer than necessary.

5. Closed-System Transfer Devices

For hazardous sterile products, such as those used in chemotherapy, closed-system transfer devices (CSTDs) are often employed. These devices mechanically prevent the transfer of environmental contaminants into the

product and the release of hazardous drug vapors or droplets into the environment. CSTDs add an extra layer of protection for both the product and the healthcare worker, and their use is increasingly mandated by regulatory standards.

THE STERILE COMPOUNDING PROCESS: A STEP-BY-STEP APPROACH

While every sterile product preparation is unique, there is a general workflow that applies to most compounding processes. This workflow is designed to maximize the safety and sterility of the final product.

Preparation and Assessment

Before any compounding begins, the pharmacist or technician reviews the prescription or medication order to ensure the drug, dose, route, diluent, and rate of administration are appropriate for the patient. Any allergies or compatibility issues are assessed. The compounding area must be certified for use, and all equipment should be functioning correctly.

Disinfection and

Setup

The workspace is thoroughly disinfected. All supplies are gathered and inspected. Vial stoppers and IV bag ports are swabbed with 70% isopropyl alcohol and allowed to dry. The personnel performing the compounding must be fully garbed, including sterile gloves that have been disinfected with alcohol.

Aseptic Manipulation

This is the core of the process. Using proper aseptic technique, the practitioner:

1. Opens sterile syringes and needles without touching the critical sites.
2. Withdraws the appropriate volume of diluent or medication from a vial, using techniques that prevent touch contamination.
3. Adds the medication to the IV bag or other final container, again maintaining sterility of all connection points.
4. Mixes the solution gently, taking care not to introduce air bubbles or create particulate matter.

Throughout these manipulations, the practitioner must be acutely aware of the first air zone and ensure that all critical sites are exposed only to clean, HEPA-filtered air. The practitioner should work at a deliberate pace, avoiding quick

QUALITY CONTROL AND can disrupt airflow or create back-contamination.

Final Check and Labeling

Once the sterile product is prepared, it is inspected for visual clarity, particulate matter, and container integrity. The final product is labeled with the patient's name, medication name, dose, diluent, administration route, expiration time, and any special instructions. A qualified pharmacist typically performs a final verification to ensure the product matches the order and that all aseptic procedures were followed correctly.

ASSURANCE IN STERILE COMPOUNDING

Quality control is not a single event but a continuous process that ensures every sterile product meets the highest standards of safety and efficacy. Key components of a comprehensive quality assurance program include:

Environmental Monitoring

Regular monitoring of the compounding environment is essential to ensure that ISO classifications are maintained. This includes:

- **Non-viable particle counting:** Measuring airborne particles to ensure HEPA filters and laminar airflow systems are functioning correctly.
- **Viable air sampling:** Testing for the presence of living microorganisms in the air using settle plates or active air samplers.
- **Surface sampling:** Swabbing work surfaces, equipment, and personnel gloves to detect microbial contamination.

- **Pressure differential monitoring:** Ensuring that positive pressure is maintained in the cleanroom relative to adjacent areas.

Media Fill Testing (Process Simulation)