
How To Stop A Period

How To Stop A Period

Downloaded from
evoluir.abar.org.br by
Autumn & Shamar

UNDERSTANDING MENSTRUATION AND THE DESIRE TO STOP A PERIOD

Menstruation is a natural biological process that signals a healthy reproductive system. Yet, for many people, monthly periods bring discomfort, inconvenience, or pain. Whether you are preparing for a special event, managing a medical condition like endometriosis, or simply seeking relief

from heavy bleeding, you may wonder **how to stop a period** temporarily or permanently. This comprehensive guide explores safe, effective methods to delay, reduce, or stop menstrual bleeding, emphasizing evidence-based advice from medical professionals.

Before considering any intervention, it is essential to understand that stopping a period should not be taken lightly. Hormonal and non-hormonal approaches carry benefits and risks. Always consult a healthcare provider before making changes to your menstrual cycle. Below,

we break down the most reliable strategies, from over-the-counter solutions to prescription medications and natural remedies.

MEDICAL METHODS TO STOP OR DELAY A PERIOD

Hormonal Birth Control: The Most Reliable Option

For those who want to **stop a period** for an extended period, hormonal contraception is the gold standard. Birth

control pills, patches, rings, and intrauterine devices (IUDs) work by stabilizing hormone levels, which prevents the uterine lining from thickening and shedding.

- **Combined oral contraceptives:**

Taking active pills continuously (skipping the placebo week) can delay bleeding indefinitely. Many women use this method for months or years under medical supervision.

- **Progestin-only pills:** Also known as the mini-pill, these can thin the uterine lining and reduce bleeding, though irregular spotting may occur initially.

- **Hormonal IUDs (e.g., Mirena):**

These devices release

levonorgestrel locally, often leading to lighter periods or complete amenorrhea within six months to a year.

- **Vaginal ring and patch:** Similar to pills, you can wear them continuously without a hormone-free break to skip periods.

Doctors typically recommend starting these methods at least a few weeks before you want to stop bleeding. It is crucial to follow the prescribed schedule to maintain effectiveness.

Nonsteroidal

Anti-Inflammatory Drugs (NSAIDs) for Heavy Bleeding

If your goal is to **stop a period** that has already started or reduce heavy flow, NSAIDs such as ibuprofen (Advil, Motrin) or naproxen (Aleve) can help. These medications reduce prostaglandins, the chemicals that trigger uterine contractions and bleeding. Taking a high dose (under a doctor's guidance) at the

onset of menstruation may decrease blood loss by 20 to 40 percent. However, NSAIDs are not a permanent solution and should not be used for extended periods due to potential stomach or kidney side effects.

Tranexamic Acid: A Prescription Option

Tranexamic acid (brand name Lysteda) is a non-hormonal medication that helps blood clots form in the uterus, reducing heavy menstrual bleeding. It is taken only during the days of heavy flow and can be effective for women who cannot

or prefer not to use hormones. Studies show it can reduce menstrual blood loss by up to 50 percent. While it does not completely **stop a period**, it can make bleeding lighter and shorter.

NATURAL AND LIFESTYLE APPROACHES TO MANAGE YOUR CYCLE

Dietary Adjustments for Lighter Flow

Some research suggests that certain nutrients and foods may influence menstrual bleeding. While natural

methods seldom stop a period entirely, they can support hormonal balance and reduce heavy flow.

- **Iron-rich foods:** Spinach, beans, red meat, and fortified cereals help maintain energy levels during menstruation but do not stop the flow.
- **Vitamin C and bioflavonoids:** Found in citrus fruits, berries, and green peppers, these compounds may strengthen blood vessels and reduce bleeding duration.
- **Ginger:** Consuming ginger tea at the start of your period may help manage pain and potentially reduce the total number of days of bleeding, according to preliminary studies.

- **Avoiding anti-clotting foods:**

Garlic, ginkgo, and high-dose vitamin E supplements can thin the blood, so limiting them during menstruation might help decrease flow.

Herbal Remedies and Traditional Practices

Many cultures use herbs to regulate menstrual cycles. However, scientific evidence is limited, and quality control varies. Always consult a doctor before using herbs, especially if you take medications.

- **Shepherd's purse (*Capsella bursa-pastoris*):** Traditionally used to reduce heavy bleeding, it may help lighten flow when taken as a tea or tincture.
- **Yarrow (*Achillea millefolium*):** Known for its astringent properties, yarrow tea is sometimes used to shorten period duration.
- **Cinnamon:** A small study suggested that cinnamon supplements could reduce menstrual bleeding and pain in some women.
- **Vitex (*Chasteberry*):** This herb is popular for balancing progesterone and may help regulate cycles over several months, though immediate effects are unlikely.

Stress Reduction and Exercise

Chronic stress can disrupt the hypothalamic-pituitary-ovarian axis, leading to irregular periods. Reducing stress through mindfulness, yoga, or meditation may help regulate your cycle. Moderate exercise improves circulation and endorphin levels, which can ease cramps and potentially reduce the duration of menstruation. However, excessive exercise can sometimes cause periods to stop altogether—a condition known as exercise-induced amenorrhea, which is not healthy for fertility or bone density.

WHEN TO CONSULT A DOCTOR ABOUT STOPPING YOUR PERIOD

It is important to differentiate between normal menstrual variation and a medical condition that requires treatment. If you experience any of the following, seek professional advice before attempting to **stop a period** on your own:

- Bleeding that lasts longer than 7 days
- Soaking through one or more pads or tampons every hour
- Passing large blood clots (larger than a quarter)
- Severe pain that interferes with daily life
- Irregular bleeding between periods
- Unexplained weight loss or fatigue

Your doctor can run tests to check for thyroid disorders, fibroids, polycystic ovary syndrome (PCOS), endometriosis, or bleeding disorders. If an underlying issue is found, addressing it may naturally regulate your cycle without needing to stop periods artificially.

RISKS AND CONSIDERATIONS OF STOPPING YOUR PERIOD

While it is generally safe to suppress menstruation using hormonal birth control, there are some potential downsides:

- **Spotting or breakthrough bleeding:** During the first few months of continuous birth control, irregular spotting is common but usually resolves.

- **Hormonal side effects:** Weight gain, mood changes, breast tenderness, or nausea may occur.
- **Increased risk of blood clots:** Combined hormonal contraceptives (containing estrogen) raise the risk of venous thromboembolism, especially in smokers or women over 35.
- **Delayed fertility return:** After stopping hormonal contraception, it may take a few months for regular cycles to resume, but there is no permanent effect on fertility.

Natural methods carry fewer systemic risks but are less reliable. Relying solely on herbs or aspirin to stop a period can lead to uncontrolled bleeding or mask a serious condition. Always prefer medical

guidance over self-experimentation.

PRACTICAL TIPS FOR MANAGING AN UNWANTED PERIOD

If you cannot stop your period completely, these strategies can help you cope:

1. **Plan ahead:** Track your cycle using an app to anticipate your period and schedule activities around it.
2. **Use period management products:** Menstrual cups, period panties, or high-absorbency tampons can provide up to 12 hours of leak-free protection.
3. **Pain relief:** Apply a heating pad to your lower abdomen, take a warm bath, or use over-the-

counter pain relievers as directed.

4. **Hydration and nutrition:** Drink plenty of water and eat balanced meals to reduce bloating and fatigue.
5. **Stay active:** Light walking or stretching can improve circulation and lessen discomfort.

THE DIFFERENCE BETWEEN STOPPING, DELAYING, AND REDUCING A PERIOD

Understanding terminology is key to setting realistic expectations:

- **Stopping a period** means achieving no bleeding at all for a defined time. This is most reliably done with continuous hormonal birth control or a hormonal IUD.

- **Delaying a period** refers to postponing the onset of bleeding, usually by taking hormones such as norethisterone (a prescription progestogen) or using extended-cycle birth control pills.
- **Reducing a period** involves making the flow lighter and shorter. NSAIDs, tranexamic acid, and lifestyle changes can contribute to this outcome.

For a one-time event like a vacation or wedding, many women ask their doctor for a short course of norethisterone, which can postpone menstruation for up to two weeks. This medication is taken three to four days before the expected period and continued until the event ends. It is not intended for long-term

use.

MYTHS ABOUT STOPPING A PERIOD

Misinformation abounds on the internet. Let's clear up a few common myths:

- **Drinking lemon juice or vinegar will stop your period.** There is no scientific evidence that acidic foods can halt menstrual flow. They may irritate your stomach.
- **Having sex will stop your period.** Orgasms can reduce cramps but do not stop bleeding. In fact, sexual activity may temporarily increase flow due to cervical stimulation.
- **You cannot get pregnant**

during your period. While unlikely, sperm can survive up to five days, and if you have a short cycle, ovulation could occur soon after menstruation ends.

- **Stopping your period is dangerous.** For most healthy women, suppressing periods with modern hormonal contraception is considered safe and does not cause toxin buildup, contrary to old myths.

CONCLUSION: MAKING AN INFORMED CHOICE

Knowing **how to stop a period** involves more than picking a quick fix—it requires understanding your body, your health needs, and the options available. The most effective and safest methods

involve medical supervision, whether through hormonal birth control, prescription medications, or lifestyle modifications. Natural remedies may offer mild support but rarely achieve complete cessation of bleeding.

If you are considering stopping your period for the long term, discuss your goals with a healthcare professional.

They can help you weigh the benefits against potential risks, monitor for side effects, and adjust the approach as needed. Remember, your menstrual cycle is a sign of reproductive health; altering it should be done with care and knowledge. By staying informed, you can take control of your cycle without compromising your wellbeing.