

Weak Convergence And Empirical Processes

Weak Convergence And Empirical Processes

Downloaded from evoluir.abar.org.br by Jillian & Becker

UNDERSTANDING WEAK CONVERGENCE AND EMPIRICAL PROCESSES

In the realm of probability theory and statistics, few concepts are as foundational—yet as subtle—as **weak convergence and empirical processes**. Whether you are a graduate student delving into asymptotic statistics, a data scientist working with large samples, or a researcher in stochastic processes, grasping these ideas is essential for making rigorous inferences from data. This article unpacks the theory of weak convergence, connects it to the behavior of empirical processes, and explores its practical implications in modern statistical analysis.

At its heart, weak convergence—also known as **convergence in distribution**—describes how a sequence of probability distributions approaches a limiting distribution. When we apply this concept to empirical processes, we obtain powerful tools for understanding the asymptotic behavior of estimators, goodness-of-fit tests, and bootstrapping methods. The marriage of weak convergence with empirical processes forms the bedrock of nonparametric statistics and functional data analysis.

What Is Weak Convergence?

Weak convergence is a type of convergence for probability measures. Formally, a sequence of random variables X_n converges weakly to X (denoted $X_n \Rightarrow X$) if the cumulative distribution functions $F_n(x)$ converge to $F(x)$ at every point x where F is continuous. This definition may seem straightforward, but its implications are far-reaching.

- **Weak convergence is weaker than almost sure convergence or convergence in probability.** It only concerns the limiting distribution, not the actual path behavior of the random variables.
- **It is central to the central limit theorem (CLT).** The CLT states that the normalized sample mean converges weakly to a normal distribution.
- **It extends naturally to stochastic processes.** For empirical processes, we consider convergence in the space of cadlag functions (right-continuous with left limits) equipped with the Skorokhod topology or the uniform topology.

The classic example of weak convergence is the convergence of the binomial distribution to the Poisson distribution for rare events, or to the normal distribution under the CLT. These results rely on the notion that the shape of the distribution stabilizes as the sample size grows.

Empirical Processes: A Bridge Between Data and Theory

An empirical process is a stochastic process constructed from a random sample. Given independent and identically distributed (i.i.d.) observations $X_1, \dots, X_n$ from a distribution P , the empirical distribution function $F_n(t)$ is defined as the proportion of observations $\leq t$. The empirical process indexed by a set of functions F is:

$$G_n(f) = \sqrt{n} (P_n f - P f)$$

where $P_n f$ is the sample average of $f(X_i)$ and $P f$ is the true expectation. The scaling by $\sqrt{n}$ ensures that the process has a non-degenerate limit.

The study of weak convergence and empirical processes investigates when G_n converges to a Gaussian process, typically a Brownian bridge. This convergence underpins many statistical procedures, including the Kolmogorov–Smirnov test, the Cramér–von Mises test, and the bootstrap.

The Glivenko–Cantelli Theorem: The Starting Point

Before diving into weak convergence, it is essential to understand the uniform consistency of the empirical distribution function. The Glivenko–Cantelli theorem states that $\|F_n - F\|_\infty \rightarrow 0$ **almost surely**. This is a strong law of large numbers for the empirical cdf, but it says nothing about the rate of convergence or the limiting distribution of the deviations.

The next natural question is: how does the scaled difference $\sqrt{n}(F_n - F)$ behave? This is precisely where weak convergence enters, leading us to the Donsker theorem.

Donsker’s Theorem: The Functional Central Limit Theorem

Donsker’s theorem is arguably the most important result linking weak convergence and empirical processes. It states that the empirical process $G_n(t) = \sqrt{n}(F_n(t) - F(t))$ converges weakly in the Skorokhod space $D[0,1]$ to a Brownian bridge $B(F(t))$, a Gaussian process with mean zero and covariance $E[B(s)B(t)] = \min(s,t) - st$.

This result provides a complete asymptotic description of the fluctuations of the empirical distribution. It enables the construction of confidence bands for the true distribution function and the derivation of asymptotic distributions for many test statistics.

- **Kolmogorov–Smirnov test:** The statistic $\sqrt{n} \|F_n - F\|_\infty$ converges to the supremum of a Brownian bridge, a distribution that is tabulated and used for goodness-of-fit testing.
- **Cramér–von Mises test:** The integrated squared difference converges to an integral of the Brownian bridge, providing a different class of tests.

Donsker’s theorem is a special case of a more general theory: the weak convergence of empirical processes indexed by classes of functions. This generalization requires conditions on the function class, such as Vapnik–Chervonenkis (VC) properties or bracketing entropy.

Key Conditions for Weak Convergence of Empirical Processes

Not every empirical process converges weakly. The key conditions revolve around the **tightness** of the sequence and the convergence of finite-dimensional distributions.

1. **Finite-dimensional convergence:** For any fixed finite set of functions $f_1, \dots, f_k$, the vector $(G_n(f_1), \dots, G_n(f_k))$ must converge in distribution to a multivariate normal. This is often ensured by the classical CLT for each fixed function.
2. **Tightness:** The sequence of processes must be stochastically equicontinuous. In practical terms, this means that small changes in the indexing function lead to small changes in the process, uniformly over the class. Tightness can be verified using entropy conditions or bracketing numbers.

When these conditions hold, we say that the empirical process is **Donsker** (or P -Donsker), meaning that G_n converges weakly to a tight Gaussian process in the space of bounded functions on F .

Applications in Modern Statistics

The framework of weak convergence and empirical processes is not merely theoretical; it drives many contemporary statistical methods.

1. BOOTSTRAP CONSISTENCY

The bootstrap, a resampling technique, relies on the convergence of the bootstrap empirical process to the same Gaussian limit as the original sample. Donsker’s theorem ensures that bootstrap confidence intervals are asymptotically valid for a wide range of statistics.

2. NONPARAMETRIC DENSITY ESTIMATION

Kernel density estimators and their functionals are often studied using empirical process theory. The weak convergence of the empirical process indexed by kernel functions leads to asymptotic normality of the integrated squared error.

3. SEMIPARAMETRIC MODELS

In models where the parameter of interest is finite-dimensional but the nuisance parameter is infinite-dimensional (e.g., Cox proportional hazards model), weak convergence of the empirical process is used to derive the asymptotic distribution of M-estimators and profile likelihoods.

4. HYPOTHESIS TESTING AND CONFIDENCE BANDS

Tests such as the Anderson–Darling test, the Shapiro–Wilk test for normality, and many others derive their critical values from the weak limit of empirical processes. Confidence bands for the entire distribution function are constructed using the quantiles of the Brownian bridge supremum.

Illustrative Example: Weak Convergence of the Empirical CDF

Suppose we sample $n = 100$ observations from a standard uniform distribution on $[0,1]$. The empirical distribution function $F_n(x)$ is a step function. For a fixed x , $F_n(x)$ is simply the proportion of sample values $\leq x$. The classical CLT tells us that $\sqrt{n}(F_n(x) - x)$ converges to a normal distribution with mean 0 and variance $x(1-x)$. That is a pointwise result.

But what about the entire function? Donsker’s theorem tells us that the whole stochastic process $\sqrt{n}(F_n(\cdot) - Id)$ converges in distribution to a Brownian bridge. This means we can make simultaneous statements about the entire cdf—for instance, that the maximum deviation over x has a known asymptotic distribution. This is far more powerful than pointwise results.

Challenges and Extensions

While the theory is elegant, real-world applications often face challenges:

- **Dependent data:** For time series or spatial data, the empirical process may converge to a different Gaussian process (e.g., a Brownian motion replaced by an Ornstein–Uhlenbeck process). Weak convergence still holds under mixing conditions, but the covariance structure changes.
- **High-dimensional settings:** When the number of functions in the class is extremely large (e.g., in machine learning), the classical Donsker conditions may fail. Modern empirical process theory in high dimensions uses tools like Rademacher complexity and concentration inequalities.
- **Non-identically distributed data:** For independent but not identically distributed observations, weak convergence of weighted empirical processes requires careful handling of the weights and the limiting covariance.

The theory continues to evolve, with recent advances in weak convergence for empirical processes indexed by unbounded functions, for functional data, and in the context of deep learning.

The Role of the Skorokhod Space and Topology

When dealing with empirical processes, the sample paths are discontinuous (step functions). The appropriate space is $D[0,1]$ or $D[0,\infty)$, the space of right-continuous functions with left limits. The Skorokhod topology is used to define weak convergence because it allows for small time shifts of jumps, which is natural when the jumps are at random sample points. In contrast, the uniform topology (supremum norm) is too strong for processes with jumps unless the limit is continuous, as

in the case of the Brownian bridge. Understanding these topologies is crucial for a rigorous treatment of weak convergence in empirical process theory.

Conclusion

Weak convergence and empirical processes together form a cornerstone of modern asymptotic statistics. From the Glivenko–Cantelli theorem to Donsker’s theorem and beyond, these concepts

provide the rigorous mathematical foundation for many statistical methods that practitioners use daily. The ability to describe the limiting behavior of sample-based processes allows us to construct valid confidence intervals, conduct hypothesis tests, and develop new methodologies with known asymptotic properties. Whether you are analyzing a simple one-sample problem or a complex semiparametric model, the language of weak convergence and empirical processes is indispensable. As data grow in size and complexity, mastering these tools becomes ever more critical for making reliable, data-driven decisions.